

LPAR1 and ITGA4 Regulate Peripheral Blood Monocyte Counts

Narelle Maugeri,^{1,2} Joseph E. Powell,¹ Peter A.C. 't Hoen,³ Eco J.C. de Geus,⁴ Gonneke Willemssen,⁴ Mathijs Kattenberg,⁴ Anjali K. Henders,¹ Leanne Wallace,¹ Brenda Penninx,⁵ Jouke-Jan Hottenga,⁴ Sarah E. Medland,¹ Viatcheslav Saviouk,⁴ Nicholas G. Martin,¹ Peter M. Visscher,¹ Gert-Jan B. van Ommen,³ Ian H. Frazer,² Dorret I. Boomsma,⁴ Grant W. Montgomery,^{1†} and Manuel A.R. Ferreira,^{1*†}

¹Queensland Institute of Medical Research, Brisbane, Australia; ²University of Queensland Diamantina Institute, Princess Alexandra Hospital, Brisbane, Australia; ³Center for Human and Clinical Genetics, Leiden University Medical Center, Leiden, The Netherlands; ⁴Netherlands Twin Register, EMGO & NCA Institute, Department of Biological Psychology, Vrije University, Amsterdam, The Netherlands; ⁵Department of Psychiatry/EMGO Institute, Vrije University Medical Center, Amsterdam, The Netherlands

Communicated by Peter J. Oefner

Received 12 April 2011; accepted revised manuscript 29 April 2011.

Published online 19 May 2011 in Wiley Online Library (www.wiley.com/humanmutation). DOI 10.1002/humu.21536

ABSTRACT: We recently mapped a quantitative trait locus for monocyte counts to chromosome 9q31 (rs7023923). Here we extend this work by showing with two independent approaches that rs7023923 regulates the expression levels of the nearby *LPAR1* gene ($P < 0.0001$), specifically implicating this gene in monocyte development. Furthermore, we tested 10 additional loci identified in the original analysis for replication in 1,122 individuals and confirm that rs6740847 near the alpha-4-integrin gene (*ITGA4*) associates with variation in monocyte counts (combined $P = 2.7 \times 10^{-10}$). This variant is in complete linkage disequilibrium ($r^2 = 1$) with a previously reported eQTL for *ITGA4* (rs2124440), indicating that this is the likely causal gene in the region. Our results indicate that rs7023923 and rs6740847 respectively upregulate *LPAR1* and downregulate *ITGA4* expression and this increases the number of monocytes circulating in the peripheral blood. Further studies that investigate the downstream mechanism involved and the impact on immune function are warranted.

Hum Mutat 32:873–876, 2011. © 2011 Wiley-Liss, Inc.

KEY WORDS: *LPAR1*; *ITGA4*; eQTL; GWAS; lysophosphatidic; *EDG2*

Genome-wide association studies (GWAS) have recently been used to identify the first genetic loci to explain a significant fraction of the variation in blood cell numbers among healthy individuals [e.g., Ferreira et al., 2009; Soranzo et al., 2009]. The expectation is that some of these loci may not only explain variation within the normal, healthy range but may also play an important role in disease, as demonstrated for a locus recently identified to regulate both eosinophil counts and asthma risk [Gudbjartsson et al., 2009]. However, despite this recent progress, the actual causal genes in many regions of confirmed association remain to be identified. Furthermore, only a small fraction of the variation between individuals can be accounted for by the identified loci, as is the case for most complex traits and diseases [Manolio et al., 2009], indicating that many more loci remain to be identified.

In a recent GWAS of hematology traits [Ferreira et al., 2009], we identified 11 loci with strong association ($P < 10^{-5}$) with monocyte counts, including a region on chromosome 9q31 that reached genome-wide significance (rs7023923, $P = 8.9 \times 10^{-14}$) and replicated unambiguously ($P = 0.001$) in an independent panel. However, we did not identify the most likely causal gene underlying the 9q31 association nor did we attempt to follow-up the 10 remaining loci that did not reach genome-wide significance. In the present study, we report results from gene-expression experiments and association analyses conducted in additional samples that identify (1) the likely causal gene in the

Additional Supporting Information may be found in the online version of this article.

[†]These authors contributed equally to this work and should be considered joint last authors.

*Correspondence to: Manuel A.R. Ferreira, Queensland Institute of Medical Research, Brisbane, Australia. E-mail: manuel.ferreira@qimr.edu.au

Contract grant sponsor: Australian National Health and Medical Research Council (NHMRC); Contract grant numbers: 241944; 339462; 389927; 389875; 389891; 389892; 389938; 443036; 442915; 442981; 496739; 552485; 552498; 613627; Contract grant sponsor: Australian Research Council; Contract grant numbers: A7960034; A79906588; A79801419; DP0212016; DP0343921; Contract grant sponsor: The Netherlands Organization for Scientific Research; Contract grant number: NWO: MagW/ZonMW; Contract grant sponsor: Twin family database for behavior genomics studies; Contract grant number: 480-04-004; Contract grant sponsor: Genotype/phenotype database for behavior genetic and genetic epidemiological studies; Contract grant number: 40-0056-98-9032;

Contract grant sponsor: Spinozapremie; Contract grant number: SPI 56-464-14192; Contract grant sponsor: CMSB: Center for Medical Systems Biology (NWO Genomics); Contract grant number: NBIC/BioAssist/RK/2008.024; Contract grant sponsors: BBMRI-NL: Biobanking and Biomolecular Resources Research Infrastructure; The VU University; Institute for Health and Care Research (EMGO+); Neuroscience Campus Amsterdam (NCA); The European Science Foundation (ESF); Genomewide analyses of European twin and population cohorts; Contract grant number: EU/QLRT-2001-01254; Contract grant sponsor: European Community's Seventh Framework Program; Contract grant number: FP7/2007-2013; Contract grant sponsor: ENGAGE; Contract grant number: HEALTH-F4-2007-201413; Contract grant sponsor: The European Science Council (ERC) Genetics of Mental Illness; Contract grant number: 230374; Contract grant sponsor: Collaborative study of the genetics of DZ twinning; Contract grant number: NIH R01D0042157-01A; Contract grant sponsors: The Genetic Association Information Network, a public-private partnership between the NIH and Pfizer Inc.; Affymetrix Inc.; Abbott Laboratories.

9q31 region and (2) a novel confirmed locus regulating monocyte counts in the general population.

We first sought to identify the causal gene underlying our previous reported association between monocyte counts and chromosome 9q31 [Ferreira et al., 2009]. Three genes are located within 250 kb of rs7023923, namely, the lysophosphatidic acid receptor 1 (*LPAR1*; MIM# 602282, also known as *EDG2*, 74 kb distant from rs7023923), olfactory receptor family 2 subfamily K (*OR2K2*, 114 kb) and *KIAA0368* (147 kb). We hypothesized that the SNP previously shown to associate with variation in monocyte counts (rs7023923) may also associate with variation in the expression levels of one of these three genes and this would be the most likely causal gene in the region.

To test this hypothesis, we analyzed gene-expression data obtained from whole blood with Illumina HumanHT-12 arrays for 48 pairs of monozygotic (MZ) twins as described in the Supp.

Table 1. Association Between the Expression Levels of *LPAR1* Measured in Whole Blood with Illumina Expression Arrays^a and rs7023923 Genotype, a SNP Previously Confirmed to Associate with Monocyte Counts [Ferreira et al., 2009]

Genotype	N	Mean	SD	Effect ^b	SE	$h^2(\%)^c$	P-value
CC	10	21.8	7.3				
CT	25	28.7	14.4	13.8	2.8	35.0	9.5×10^{-6}
TT	13	48.6	13.3				

^aExpression of *LPAR1* was assessed by probe ILMN_1701441 that is present in the Illumina HumanHT-12 v3.0 Beadchip used.

^bEffect corresponds to the slope from a linear regression of expression levels on the number of T alleles. Sex was not a significant predictor of the expression of the transcripts tested and so was not included as a covariate.

^c h^2 represents the proportion of the between-individual variation in gene expression explained by the SNP.

N, sample size; SE, standard error.

Methods. After quality control, expression levels were available for transcripts ILMN_1701441 in *LPAR1*, ILMN_1745801 in *OR2K2* and ILMN_1847822 in *KIAA0368* (Supp. Table S1). Genotype data for rs7023923 were also available for each twin pair as described previously [Ferreira et al., 2009]. Linear regression considering additive single nucleotide polymorphism (SNP) effects was then used to test whether rs7023923 was associated with the expression of each of the three transcripts, considering the mean expression for each MZ pair. SNP rs7023923 was found to be a significant predictor of the expression levels of *LPAR1* ($P = 9.5 \times 10^{-6}$; Table 1) but not of *OR2K2* or *KIAA0368* ($P > 0.05$). This SNP explained 35% of the between-individual variation in *LPAR1* expression; the rs7023923T allele that associated with increased *LPAR1* expression also associated with increased monocyte counts. We replicated these results using a different technology (DeepSAGE, Supp. Methods) in 94 independent samples ($P = 5.4 \times 10^{-5}$, Supp. Fig. S1). Furthermore, results from a large study of the monocyte transcriptome [Zeller et al., 2010] strongly support our observation (Supp. Table S2). Thus, our results specifically implicate *LPAR1* in the regulation of monocyte counts.

Lysophosphatidic acid (LPA) is a ubiquitous extracellular growth factor produced from membrane phospholipids that exerts physiological effects through five high-affinity cognate receptors [Choi et al., 2010]. The first to be identified was the LPA1 receptor [Hecht et al., 1996], which is encoded by the *LPAR1* gene that we now implicate in monocyte development. This receptor is moderately expressed in many human tissues, including monocytes [Choi et al., 2010; Goetzl et al., 2000], and it has been suggested to mediate the capacity of LPA to suppress cellular apoptosis [Goetzl et al., 1999]. Together with our results, these data suggest that the rs7023923T allele increases the numbers of peripheral blood monocytes by increasing the




Figure 1. Regional plot of association between SNPs in chromosome 2q31 and monocyte counts. Results are shown for the combined analysis of the original GWAS panel ($N = 4,225$) and the new replication panel ($N = 1,122$) and include the most associated SNP (rs6740847, shown in blue, $P = 2.7 \times 10^{-10}$) as well as all available SNPs within 150 kb of rs6740847. The color of the remaining markers reflects the linkage disequilibrium (r^2) with rs6740847 (increasing red hue associated with increasing r^2). The recombination rate (second y axis) is plotted in light blue and is based on the CEU HapMap population. Exons for each gene are represented by vertical bars, based on all isoforms available from the March 2006 UCSC Genome Browser assembly.

surface expression of the LPA1 receptor, which in turn upregulates LPA signaling pathways that promote monocyte proliferation or survival.

Next, we followed-up the 10 top loci that did not reach genome-wide significance in our original GWAS of monocyte counts in a further 1,122 unrelated individuals ascertained from the general population and tested as described in the Supp. Methods. Of those 10 SNPs, only one located in chromosome 2q31 showed a convincing association with monocyte counts in the replication panel (rs6740847, uncorrected $P = 0.002$; Supp. Table S3). In the combined analysis of the original GWAS and replication panels ($N = 5,347$) the association with this locus was highly significant ($P = 2.7 \times 10^{-10}$). Thus, our results indicate that a novel locus that regulates monocyte counts is located on chromosome 2q31.

The rs6740847 variant is located 13 kb upstream from the transcription start site of the alpha-4-integrin gene (*ITGA4*; MIM# 192975; Fig. 1). We therefore tested whether rs6740847 was a significant predictor of *ITGA4* expression levels in the blood, but found no significant association with the specific probe for this gene available in the Illumina expression array used (Supp. Table S4) or for the nearby *CERKL* (MIM# 608381) gene (four probes, all with $P > 0.05$). We sought to confirm these negative results using deepSAGE analysis in 94 independent samples, but instead observed that the rs6740847G allele was associated with decreased *ITGA4* expression ($P = 0.02$; Supp. Fig. S2). No association was observed with *CERKL* (not shown). Furthermore, we subsequently noted that rs6740847 was in complete LD ($r^2 = 1$) with an eQTL for *ITGA4* in monocytes (rs2124440, $P = 2.2 \times 10^{-33}$; Supp. Table S5) recently identified by Zeller et al. [2010]. These results thus suggest that *ITGA4* is indeed the likely causal gene underlying the rs6740847 association with monocyte counts. The negative results observed in our microarray eQTL analysis likely reflect the low power provided by the analysis of 48 samples (57% for $\alpha = 0.05$).

Alpha-4-integrin (or very late activation antigen-4, VLA-4) is expressed on the cell surface of most circulating and tissue-resident mononuclear leukocytes, including monocytes [Hemler et al., 1990]. It preferentially binds VCAM-1 and fibronectin produced by stromal cells and this adhesive interaction is crucial for leukocyte homing and trafficking through the bloodstream [Kukreti et al., 1997; Oostendorp and Dörmer, 1996]. Indeed, blocking alpha-4-integrin [Issekutz, 1991; Papayannopoulou and Nakamoto, 1993] or disrupting VCAM-1 expression [Gurtner et al., 1995] led to a significant increase in the number of circulating leukocytes. These experimental observations are consistent with our results for rs6740847, which suggest that the G allele increases monocyte counts in the peripheral blood by decreasing *ITGA4* expression.

In summary, results from our analyses specifically implicate the *LPAR1* and *ITGA4* genes in the regulation of monocyte counts, with the two variants identified (rs7023923 and rs6740847) jointly accounting for 1.2% ($P = 0.0015$) of the between-individual variation in the replication cohort tested. Further studies are warranted that investigate the downstream molecular mechanisms involved and the potential impact of monocyte numbers in inflammation and the innate immune response.

Acknowledgments

We thank the Brisbane twins and their families for their participation, as well as Dixie Statham, Ann Eldridge, Marlene Grace; Lisa Bowdler, Steven Crooks, Megan Campbell, Kalpana Patel, and Anthony Caracella; David Smyth and Harry Beeby.

References

- Aitken JF, Green AC, MacLennan R, Youl P, Martin NG. 1996. The Queensland Familial Melanoma Project: study design and characteristics of participants. *Melanoma Res* 6:155–165.
- Choi JW, Herr DR, Noguchi K, Yung YC, Lee CW, Mutoh T, Lin ME, Teo ST, Park KE, Mosley AN, Chun J. 2010. LPA receptors: subtypes and biological actions. *Annu Rev Pharmacol Toxicol* 50:157–186.
- Ferreira MA, Hottenga JJ, Warrington NM, Medland SE, Willemsen G, Lawrence RW, Gordon S, de Geus EJ, Henders AK, Smit JH, Campbell MJ, Wallace L, Evans DM, Wright MJ, Nyholt DR, James AL, Beilby JP, Penninx BW, Palmer LJ, Frazer IH, Montgomery GW, Martin NG, Boomsma DI. 2009. Sequence variants in three loci influence monocyte counts and erythrocyte volume. *Am J Hum Genet* 85:745–749.
- Freeman MF, Tukey JW. 1950. Transformations related to the angular and the square root. *Ann Math Stat* 21:607–611.
- Goetzl EJ, Kong Y, Mei B. 1999. Lysophosphatidic acid and sphingosine 1-phosphate protection of T cells from apoptosis in association with suppression of Bax. *J Immunol* 162:2049–2056.
- Goetzl EJ, Kong Y, Voice JK. 1999. Cutting edge: differential constitutive expression of functional receptors for lysophosphatidic acid by human blood lymphocytes. *J Immunol* 164:4996–4999.
- Gudbjartsson DF, Bjornsdottir US, Halapi E, Helgadóttir A, Sulem P, Jonsdóttir GM, Thorleifsson G, Helgadóttir H, Steinthorsdóttir V, Stefansson H, Williams C, Hui J, Beilby J, Warrington NM, James A, Palmer LJ, Koppelman GH, Heinzmann A, Krueger M, Boezen HM, Wheatley A, Altmüller J, Shin HD, Uh ST, Cheong HS, Jonsdóttir B, Gislason D, Park CS, Rasmussen LM, Porsbjerg C, Hansen JW, Backer V, Werge T, Janson C, Jönsson UB, Ng MC, Chan J, So WY, Ma R, Shah SH, Granger CB, Quyyumi AA, Levey AI, Vaccarino V, Reilly MP, Rader DJ, Williams MJ, van Rij AM, Jones GT, Trabetti E, Malerba G, Pignatti PF, Boner A, Pescollenderung L, Girelli D, Olivieri O, Martinelli N, Ludviksson BR, Ludviksdóttir D, Eyjolfsson GI, Arnar D, Thorgeirsson G, Deichmann K, Thompson PJ, Wjst M, Hall IP, Postma DS, Gislason T, Gulcher J, Kong A, Jonsdóttir I, Thorsteinsdóttir U, Stefansson K. 2009. Sequence variants affecting eosinophil numbers associate with asthma and myocardial infarction. *Nat Genet* 41:342–347.
- Gurtner GC, Davis V, Li H, McCoy MJ, Sharpe A, Cybulsky MI. 1995. Targeted disruption of the murine VCAM1 gene: essential role of VCAM-1 in chorioallantoic fusion and placentation. *Genes Dev* 9:1–14.
- Hecht JH, Weiner JA, Post SR, Chun J. 1996. Ventricular zone gene-1 (vzg-1) encodes a lysophosphatidic acid receptor expressed in neurogenic regions of the developing cerebral cortex. *J Cell Biol* 135:1071–1083.
- Hemler ME, Elices MJ, Parker C, Takada Y. 1990. Structure of the integrin VLA-4 and its cell–cell and cell–matrix adhesion functions. *Immunol Rev* 114:45–65.
- Hestand MS, Klingenhoff A, Scherf M, Ariyurek Y, Ramos Y, van Workum W, Suzuki M, Werner T, van Ommen GJ, den Dunnen JT, Harbers M, 't Hoen PA. 2010. Tissue-specific transcript annotation and expression profiling with complementary next-generation sequencing technologies. *Nucleic Acids Res* 38:e165.
- Howie BN, Donnelly P, Marchini J. 2009. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet* 5:e1000529.
- Issekutz TB. 1991. Inhibition of in vivo lymphocyte migration to inflammation and homing to lymphoid tissues by the TA-2 monoclonal antibody. A likely role for VLA-4 in vivo. *J Immunol* 147:4178–4184.
- Kukreti S, Konstantopoulos K, Smith CW, McIntire LV. 1997. Molecular mechanisms of monocyte adhesion to interleukin-1beta-stimulated endothelial cells under physiologic flow conditions. *Blood* 89:4104–4111.
- Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorf LA, Hunter DJ, McCarthy MI, Ramos EM, Cardon LR, Chakravarti A, Cho JH, Guttacher AE, Kong A, Kruglyak L, Mardis E, Rotimi CN, Slatkin M, Valle D, Whittemore AS, Boehnke M, Clark AG, Eichler EE, Gibson G, Haines JL, Mackay TF, McCarroll SA, Visscher PM. 2009. Finding the missing heritability of complex diseases. *Nature* 461:747–753.
- McGregor B, Pitzner J, Zhu G, Grace M, Eldridge A, Pearson J, Mayne C, Aitken JF, Green AC, Martin NG. 1999. Genetic and environmental contributions to size, color, shape, and other characteristics of melanocytic naevi in a sample of adolescent twins. *Genet Epidemiol* 16:40–53.
- Oostendorp RA, Dörmer P. 1996. VLA-4-mediated interactions between normal human hematopoietic progenitors and stromal cells. *Leuk Lymphoma* 24:423–435.
- Papayannopoulou T, Nakamoto B. 1993. Peripheralization of hemopoietic progenitors in primates treated with anti-VLA4 integrin. *Proc Natl Acad Sci USA* 90:9374–9378.
- Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC. 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 81:559–575.

- Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26:139–26:140.
- Soranzo N, Spector TD, Mangino M, Kühnel B, Rendon A, Teumer A, Willenborg C, Wright B, Chen L, Li M, Salo P, Voight BF, Burns P, Laskowski RA, Xue Y, Menzel S, Altshuler D, Bradley JR, Bumpstead S, Burnett MS, Devaney J, Döring A, Elosua R, Epstein SE, Erber W, Falchi M, Garner SE, Ghorji MJ, Goodall AH, Gwilliam R, Hakonarson HH, Hall AS, Hammond N, Hengstenberg C, Illig T, König IR, Knouff CW, McPherson R, Melander O, Mooser V, Nauck M, Nieminen MS, O'Donnell CJ, Peltonen L, Potter SC, Prokisch H, Rader DJ, Rice CM, Roberts R, Salomaa V, Sambrook J, Schreiber S, Schunkert H, Schwartz SM, Serbanovic-Canic J, Sinisalo J, Siscovick DS, Stark K, Surakka I, Stephens J, Thompson JR, Völker U, Völzke H, Watkins NA, Wells GA, Wichmann HE, Van Heel DA, Tyler-Smith C, Thein SL, Kathiresan S, Perola M, Reilly MP, Stewart AF, Erdmann J, Samani NJ, Meisinger C, Greinacher A, Deloukas P, Ouwehand WH, Gieger C. 2009. A genome-wide meta-analysis identifies 22 loci associated with eight hematological parameters in the HaemGen consortium. *Nat Genet* 41:1182–1190.
- Willemsen G, de Geus EJ, Bartels M, van Beijsterveldt CE, Brooks AI, Estourgie-van Burk GF, Fugman DA, Hoekstra C, Hottenga JJ, Klufft K, Meijer P, Montgomery GW, Rizzu P, Sondervan D, Smit AB, Spijker S, Suchiman HE, Tischfield JA, Lehner T, Slagboom PE, Boomsma DI. 2010. The Netherlands Twin Register biobank: a resource for genetic epidemiological studies. *Twin Res Hum Genet* 13:231–245.
- Zeller T, Wild P, Szymczak S, Rotival M, Schillert A, Castagne R, Maouche S, Germain M, Lackner K, Rossmann H, Eleftheriadis M, Sinning CR, Schnabel RB, Lubos E, Mennerich D, Rust W, Perret C, Proust C, Nicaud V, Loscalzo J, Hübner N, Tregouet D, Münzel T, Ziegler A, Tiret L, Blankenberg S, Cambien F. 2010. Genetics and beyond—the transcriptome of human monocytes and disease susceptibility. *PLoS One* 5:e10693.