

<b>ECG parameter</b>	<b>Rate</b>	<b>MZ correlation</b>	<b>DZ correlation</b>
<b>TpTe</b>	<b>Low</b>	0.58 (0.35-0.72)	0.01 (-0.32-0.33)
	<b>Medium</b>	0.67 (0.51-0.78)	0.11 (-0.27-0.44)
	<b>High</b>	0.63 (0.44-0.75)	-0.18 (-0.57-0.36)
	<b>Resting</b>	0.56 (0.36-0.70)	0.08 (-0.25-0.37)
<b>Th</b>	<b>Low</b>	0.58 (0.34-0.74)	0.38 (0.08-0.59)
	<b>Medium</b>	0.56 (0.33-0.71)	0.43 (0.15-0.62)
	<b>High</b>	0.54 (0.29-0.71)	0.39 (0.11-0.61)
	<b>Resting</b>	0.45 (0.19-0.63)	0.19 (-0.06-0.41)
<b>QT</b>	<b>Low</b>	0.69 (0.52-0.80)	0.07 (-0.30-0.41)
	<b>Medium</b>	0.57 (0.37-0.71)	0.24 (-0.07-0.50)
	<b>High</b>	0.47 (0.20-0.65)	0.25 (-0.09-0.52)
	<b>Resting</b>	0.52 (0.29-0.68)	0.09 (-0.17-0.34)
<b>QRS</b>	<b>Low</b>	0.46 (0.20-0.64)	0.10 (-0.24-0.41)
	<b>Medium</b>	0.49 (0.25-0.66)	0.08 (-0.25-0.39)
	<b>High</b>	0.39 (0.14-0.59)	0.25 (-0.24-0.58)
	<b>Resting</b>	0.30 (0.07-0.50)	0.00 (-0.29-0.29)

**Supplemental Table 2:** Monozygotic and dizygotic twin correlations from the saturated model including participants taking cardioactive medication. 99% CIs are shown in parentheses