

The Familial Clustering of Age at Menarche in Extended Twin Families

Stéphanie M. van den Berg · Dorret I. Boomsma

Received: 27 September 2006 / Accepted: 7 May 2007 / Published online: 31 May 2007
© Springer Science+Business Media, LLC 2007

Abstract The timing of puberty is complex, possibly involving many genetic factors that may interact with the gradual increase in gonadotropin-releasing hormone (GnRH) pulse frequency at the beginning of puberty is environmental influences. Familial resemblance for age at menarche was studied in a sample of 4,995 female twins, 1,296 sisters, 2,946 mothers and 635 female spouses of male twins. They had indicated their age at menarche and it is likely that genetic factors interact with part of a larger longitudinal survey. We assessed assortative mating for age at menarche, gene-environment interaction and estimated the heritability of individual differences in the timing of pubertal development effects and estimated the heritability of individual differences in pubertal timing. There was significant evidence of gene-environment interaction, accounting for 1.5% of the variance. There was no indication of consistent mate assortment on age at menarche. Individual differences in age at menarche are highly heritable, with additive genetic factors explaining at least 70% of the true variation. An additional 1.5% of the variation can be explained by a genotype-environment interaction effect where environmental factors are more important in individuals genetically predisposed for late menarche.

Keywords Menarche · Heritability · Gene-environment interaction · Assortative mating

Introduction

Puberty refers to the re-activation of the hypothalamic-pituitary-gonadal axis, culminating in sexual maturation

had their first menses, application of a multifactorial threshold model resulted in an estimate of 30% for the heritability of menarche before age 12 (van den Berg et al. 2006). Heritability studies on age at menarche, most of them retrospective and only using adult twin samples, usually find estimates of about 50% or higher (Towne et al. 2005). When controlling for measurement error, Pickles et al. (Pickles et al. 1998) found that nearly all true variance in age at menarche could be attributed to additive genetic effects.

A drawback of these studies is that they did not go beyond the application of a relatively simple genetic model, that is, they examined neither assortative mating nor genotype-environment interaction. If there are interactions between genotype and environmental factors that are either shared or non-shared between family members, the heritability coefficient is biased when these interactions are

Edited by Stacey Cherny

S. M. van den Berg (✉) · D. I. Boomsma
Department of Biological Psychology, VU University
Amsterdam, Van der Boerhorststraat 1, P.O. Box 1081 BT,
Amsterdam, The Netherlands
e-mail: SM.van.den.Berg@psy.vu.nl

ignored. For instance, the variance attributable to an interTwin Registry (NTR). In 1991, the NTR started a longitudinal survey study of health, personality and lifestyle. environmental effects, when ignored, is attributed to non-Surveys were sent out in 1991, 1993, 1995, 1997, 2000 and shared environmental factors, while it might make more2002 to adolescent and adult twins and their family sense to attribute this variance to genetic factors (Falconer and Mackay1996). Interaction between genetic effects and waves, siblings were included since 1995. Parents of twins non-shared environmental effects can be investigated by were asked to participate in 1991, 1993, 1995 and 2002 and correlating the difference between the two individuals fromspouses (married or unmarried) since 2000. Families with a monozygotic (MZ) twin pair with the average age atadolescent and young adult twins were recruited through menarche (or, equivalently, the sum of the ages) of that twinCity Councils in 1990–1991 and in 1992–1993. After 1993 pair (Jinks and Fulker1970). With a positive correlation, an effort was also made to recruit adult and older twins the relative impact of the environment is larger with indi- through a variety of approaches. Further details on re- individuals with a familial predisposition for late menarche. sponse rates, response bias and demographic characteristics

Another commonly made but untested assumption in if of the sample can be found elsewhere (Boomsma et al. studies estimating heritability is that mating is random with2002, Stubbe et al.2005, Vink et al. 2004, Koopmans regard to the phenotype of interest: one assumes that matesal. 1999). Data from female spouses and first-degree are not genetically correlated regarding age at menarche. female relatives of male twins were used to examine Of course, males do not have menses, but they are never assortative mating. For the genetic model fitting, data from theless carriers of the genes that influence the age at male-female twin pairs, female twins from opposite sex menarche in their female offspring. If males from families twins, female full siblings and mothers were used; data with precocious females have a tendency to mate with from half-siblings and multiples other than twins were relatively precocious females, and males from familiesexcluded from the analyses. We had data on 1,340 families with late puberty onset choose to mate with relatively late with MZ female twins, 793 families with DZ female twins, females, the offspring in one family is expected to be more1,078 families with DZ unlike sex twins and 1,141 families genetically similar with respect to age at menarche than without female twins or where zygosity was unknown (16 under random mating (Falconer and Mackay1996). The families). We had data on 4,995 individual twins, 1,296 assortment need not take place based on the phenotype sisters, 2,946 mothers and 635 female spouses of male se: a genetic correlation is also induced when assortment twins. Average family size (excluding spouses) was 2.1 takes place on a phenotype genetically correlated with age (SD = 1.0; median = 2, mode = 1, maximum = 7). at menarche, for example BMI (Kaprio et al.1995). For 993 female twin pairs zygosity was based on DNA

An increase in the genetic similarity of the offspring has polymorphisms. For the other same sex twin pairs, zygosity important consequences for quantitative genetic modeling was based on eight items on physical similarity and the Usually, it is assumed that the additive genetic effects frequency of confusion of the twins by parents, other affecting a phenotype correlate 1/2 in full sisters and family members and strangers. Agreement between female dizygotic (DZ) twins. Under positive phenotypic zygosity based on these items and zygosity based on DNA assortment (males mating with females with puberty timing was 98% (Willemsen et al.2005). Average age at the time similar to that of their mothers and sisters), the genetic of the first available report on age at menarche was 25 correlation is larger than 1/2. When ignored, heritability is years for twins, 30 years for siblings and 46 years for underestimated. mothers.

The familial clustering of age at menarche was studied in a large sample of female twins, their mothers, their full Study design sisters and female spouses and first-degree relatives of male twins. All were asked to indicate their age at men-In 1991, 1993 and 1995 female participants were asked to arche as part of a longitudinal survey study. indicate the age at menarche in years and months. In 2000 and 2002, participants were only asked to indicate the age in years. In addition, in 1993, 1995 and 2002, participants were asked whether they had had their first menstruation prior to indicating the age at which it had occurred. All data concerning age at menarche were rescaled to number of months, where 6 months were added to the data from 2000

Subjects and methods

Subjects

Analyses were based on data from multiples and their and 2002. This way we avoided bias due to the fact that we family members who are registered with the Netherlands only had data on the age in years.

Data-analysis

Over the years, the reports on menarche were not entirely consistent. Data were discarded when the reported age at menarche was higher than the age at the time of the questionnaire and when one of more than two reports deviated more than 12 months from the other reports when these were more consistent. Data points were also ignored when participants indicated they had not yet had their menses but nevertheless reported an age. Total discarded data was less than 1%.

Age at menarche was correlated between spouses of male twins and first-degree relatives of the male twins were correlated to test for genotype-environment interaction. Sums and differences in age at menarche in MZ twins were assessed and subsequently modeled to estimate heritability in a quantitative genetic analysis. Model fit was assessed by comparing the fit of nested models using likelihood ratio tests. The probability for a Type I error was fixed at 5% for each test.

Results

Consistency

The within-person correlations over time ranged from .71 to .89. The first available report correlated .97 with the average reported age at menarche. The first available report was considered to be the most reliable and used for the analyses. Average age at menarche was 161.7 months (13.5 years, SD = 17.0, minimum = 96, maximum = 246).

Assortative mating

The correlation between the age at menarche in female spouses of male twins is $\beta = .08$, 95% CI $\beta = .02, .14$, was significant ($r = .17, p = .001$), and similarly for twin 2 difference ($r = .14, p = .006$). Taken together, these results suggest that in 2002, the environmental variance factors that are shared in (male) twins. Correlations between age at menarche of mothers and that of their spouses of their male twin offspring was estimated at .08 ($N = 345, \beta = .02, .18$). Correlations between spouses of twins and sisters was $\beta = .14$ and just significant ($N = 229, \beta = .06, \beta = .005$) and the correlation between a male twin spouse and his female co-twin was $\beta = .08$ ($N = 165, \beta = .02, .14$). The inconsistent pattern of correlations between spouses and first-degree relatives is negative for siblings and positive for mothers and no correlation between the female spouses of twins suggests no assortment for age at menarche or traits related to it. Therefore in the genetic component was ignored in further analyses.

modeling, we assumed that mating is random in the population with regards to age at menarche and that the correlation between any additive genetic effects in female offspring is 1/2.

Gene-environment interaction

There was a significant correlation between the sums and absolute differences in MZ twins, $r = .13, N = 1,122, p < .01$, suggesting an interaction between shared genetic or environmental effects and non-shared environmental effects (Jinks and Fulker 1990). The positive sign of the correlation indicates that non-shared environmental effects seem to have more impact in twins that have a familial predisposition for late menarche. Of course, these non-shared environmental effects might simply reflect error variance: possibly, there is more unreliability in the reports of females with late menarche. When analyzing the MZ twin data from each measurement separately, we see the lowest correlation between MZ sums and absolute differences in 1991 ($\beta = .01, N = 346$) and the highest in 2002 ($\beta = .24, N = 577$). The variance of the absolute difference scores in 2002 is 144, while the within-twin absolute differences for 2002 and 2000 reports have variances of 59 and 48, for twin 1 and twin 2, respectively. Thus, the larger part of the environmental variance seems to be true environmental variance, rather than measurement unreliability.

In 2002, we found no correlation between reported age at menarche and absolute difference between the 2000 and 2002 reports for the first twin ($r = .07, N = 513, p = .13$), but a significant correlation for the second twin ($r = .16, N = 524, p < .001$). The 2002 twin 1 + twin 2 summed report correlated neither with the within-twin 1 differences, $r = .07, p = .19$, nor with the within-twin 2 differences, ($r = .08, p = .09$). But again, the correlation between the absolute difference between

2002 and 2000 difference between twin 1 and twin 2 difference ($r = .14, p = .006$). Taken together, these results suggest that in 2002, the environmental variance factors that are shared in (male) twins. Correlations between age at menarche of mothers and that of their spouses of their male twin offspring was estimated at .08 ($N = 345, \beta = .02, .18$). Correlations between spouses of twins and sisters was $\beta = .14$ and just significant ($N = 229, \beta = .06, \beta = .005$) and the correlation between a male twin spouse and his female co-twin was $\beta = .08$ ($N = 165, \beta = .02, .14$).

It should be noted that the GE interaction reflected by the correlation between sum and differences is small, explaining only 1.5% of the variation in the reports. Whatever the extent to which this genotype-dependent environmental variance reflects true variance, this interaction component was ignored in further analyses.

Analysis of means, variances and covariances

A base model was constructed specifying for each of the four different types of families three mean values and three variances for twins, siblings and mother. In addition for each group four covariances were specified for mother–daughter, sibling–sibling, sibling–twin and twin–twin relationships (see Table 1). The twin–twin covariance was not estimated for the opposite sex twin group, nor was the twin–sibling covariance in the male twins/unknown zygosity group as there were no data for this relationship. The $-2LL$ χ^2 statistic was 77233.67 with 9,195 degrees of freedom. In consecutive steps, mean values, variances and covariances were equated across groups, and also within groups, leading to a reference model where all mean values were equal, twin variances were equal, sibling variances were equal and mother variances were equal across twins, siblings and mothers could be equated without significant loss of fit. The correlation between age at menarche and age at the time of report was only .06. The correlation between birth year and reported age at menarche was $-.06$. The distribution in twins and siblings had however significant excess kurtosis, being more leptokurtotic than expected under the assumption of normality. This was not observed in the mothers where the distributions for old and young mothers were slightly platykurtotic but not significantly so.

sibling reports were not available from the 1991 and 1993 surveys and were therefore more often than for twins based on reports from the 2000 and 2002 surveys, where age at menarche was only reported in years. This lack of precision might well have increased measurement error. Indeed, in the twin data for example, the variance in 2000 and 2002 was 307 and 312, respectively, whereas in the preceding waves the variance ranged between 222 and 251. Moreover, MZ twin correlations were .60 and .57 in 2000 and 2002, and between .81 and .87 in the preceding waves. In further analyses, therefore, it was assumed that the surplus sibling and mother variances were due to extra measurement variance. There was no evidence for a systematic telescoping effect where the age at menarche is reported more forward or more backward in time with increasing duration between event and report: means of twins, siblings and mothers could be equated without significant loss of fit. The correlation between age at menarche and age at the time of report was only .06. The correlation between birth year and reported age at menarche was $-.06$. The distribution in twins and siblings had however significant excess kurtosis, being more leptokurtotic than expected under the assumption of normality. This was not observed in the mothers where the distributions for old and young mothers were slightly platykurtotic but not significantly so.

Table 1 Observed covariance (below diagonal) and correlation matrices (above diagonal) for the different types of families (families with unknown twin zygosity not presented)

	Twin 1	Twin 2	Sibling 1	Sibling 2	Mother
MZ twins					
Twin 1	277.46	.71	.39	.39	.33
Twin 2	195.82	277.46	.39	.39	.33
Sibling 1	121.02	121.02	349.38	.37	.30
Sibling 2	121.02	121.02	129.11	349.38	.30
Mother	98.74	98.74	99.90	99.90	323.22
DZ twins					
Twin 1	267.04	.30	.26	.26	.32
Twin 2	79.24	267.04	.26	.26	.32
Sibling 1	68.28	68.28	249.46	.09	.31
Sibling 2	68.28	68.28	23.34	249.46	.31
Mother	90.94	90.94	86.97	86.97	305.74
Unlike sex twins					
Twin 1	264.56	Ø	.31	.31	.31
Twin 2	Ø	264.56	.31	.31	.31
Sibling 1	85.78	85.78	289.88	.33	.33
Sibling 2	85.78	85.78	96.37	289.88	.33
Mother	90.16	90.16	99.11	99.11	313.20

The sibling–sibling covariance, twin–sibling covariance and DZ twin covariance could all be equated. These were however significantly smaller than the MZ twin covariance, $\chi^2(1) = 202.16$, suggesting a genetic component in the individual differences. The reference model had a $-2LL$ χ^2 statistic of 77267.43 with 9,229 degrees of freedom. The covariance structure estimated using this reference model is given in Table 2. The DZ twin/sibling covariance is a little less than half the MZ twin/sibling covariance, but the mother–daughter covariance is practically half the MZ covariance; taken together this suggests no non-additive gene action.

To put this interpretation to the test, a genetic model was fitted using maximum likelihood estimation on the raw data, hypothesizing additive genetic variance (A), dominance genetic variance (D) and non-shared environmental variance (E), in addition to surplus variance in siblings (S) and mothers (M). The expectation for twin variance was $A + D + E$, for siblings, $A + D + E + S$ and for mothers $A + D + E + M$. Since MZ twins are genetically identical, the expectation for the MZ twin covariance was $A + D$. For DZ twin and siblings covariances the expectation was $1/2A + 1/4D$, and for the mother–daughter covariance $1/2A$, since parents only transmit additive genetic variance. The model had a $-2LL$ χ^2 statistic of 77268.09 with 9,230 degrees of freedom. The estimates for A, D and E were 189.42, 0.65 and 82.02, respectively, with D not

Table 2 The expected covariance structure based on the reference model (top) and the genetic model (bottom)

	Twin 1	Twin 2	Sibling 1	Sibling 2	Mother
Twin 1	271.70	.70/.33	.32	.32	.33
Twin 2	MZ: 190.07/DZ: 90.27	271.70	.32	.32	.33
Sibling 1	90.27	90.27	295.41	.31	.31
Sibling 2	90.27	90.27	90.27	295.41	.31
Mother	95.67	95.67	95.67	95.67	316.75
	Twin 1	Twin 2	Sibling 1	Sibling 2	Mother
Twin 1	272.09	.70/.35	.33	.33	.32
Twin 2	MZ: 190.06/DZ: 94.87	272.09	.33	.33	.32
Sibling 1	94.87	94.87	297.81	.32	.31
Sibling 2	94.87	94.87	94.87	297.81	.31
Mother	94.71	94.71	94.71	94.71	315.77

Correlations above the diagonal, covariances below the diagonal

significantly different from 0, $P < 0.01$. S and M (e.g., maternal effects). Given the present study design it is not possible to disentangle dominant gene action, epistatic plus error variance in siblings and mothers, broad heritability is estimated at $(A + D)/(A + D + E) = 70\%$ (95% CI 67%, 72%). With the constraint $D = 0$ the results for narrow heritability were exactly the same.

Discussion

The age at menarche in sisters of male twins correlated significantly with the age at menarche of the spouses of male twins. This was a negative correlation, where a late menarche in the male twin's sisters predicted early menarche in the male twin's spouse. However at the same time, there was a significant correlation between the age at menarche in the mother and the male twin's spouse was positive. Thus, there is consistent evidence that the genetic effects that affect age at menarche are correlated in mating partners. This is important since in quantitative genetic modeling it is often assumed that this is not the case. If a positive genetic correlation between spouses truly exists but is ignored, heritability is underestimated. In the case of age at menarche it can be safely assumed that the additive genetic effects on age at menarche correlate 1/2 in sisters.

Based on the quantitative genetic modeling, significant gene-environment interaction effect found here. Late evidence of additive genetic contribution to individual differences in age at menarche was found, accounting for 70% of the variance in the reports. Non-additive genetic effects affecting steroid levels. Alternatively, late menarche might action was not detected. If there has been selection for the trait over the course of evolution, it probably was selection for an intermediate optimum.

It should be noted however, that the modeling was based on the assumption that the heritability was constant across generations. Furthermore, it was assumed that there are no environmental influences that are shared among offspring.

availability, olfactory cues, diet and stress (Sisk and Foster 2004; Ebling 2005). However, the gene-environment interaction effect was relatively small, accounting for only 1.5% of the variance in twins. Moreover, it cannot be ruled out that at least part of the GE interaction variance reflects and error variance associated with reports of late menarche. Whatever the case, the bulk of the variance seems to result from the effects of genes and environment that act independently, with genes explaining most of the variance. Our results showed an estimate of 70% for the amount of total variance accounted for by additive genetic variation. Since in the analysis gene-environment interaction was ignored, its variance is attributed to non-shared variance. One may therefore conclude that between 70 and 71.5% of the total variance can be attributed to the effects of genotype (Falconer and Mackay 1996).

Total variance however includes variance due to measurement error. Therefore the relative impact of genetic influences may be underestimated. Given the average ages of twins, siblings and the mothers, there was a long interval between the actual menarche and the time of report that may have resulted in substantial error variance. To some extent the error variance was accounted for by allowing larger variances for siblings and mothers, but not all error variance could be accounted for. Although there is indeed evidence of unreliability in retrospective menarche reports the error seems to be symmetric around the actual age at menarche since the mean values in the three groups were equal (Must et al 2002; Greif and Ulman 1982; Pillemer et al. 1987). We did however observe leptokurtic distributions in the reports. These might be indicative of a tendency to report menarche more closely to the average age than expected with a normal distribution, possibly related to a memory bias towards the population mean. A leptokurtic distribution for recalled age at menarche has been observed in other studies (Treloar and Martin 1990; Zacharias et al 1970). However the present results should be interpreted with care, since the test is highly sensitive to the assumption of a continuous variable, which is not the case here where in two surveys one could indicate age at menarche only in years.

We found no systematic telescoping effect, where remembered events are placed more forward or more backward in time with increasing time between actual event and recall (Pickles et al 1998). However, it should be noted such an effect is confounded with the generally observed decrease in age of puberty onset (Wyshak and Frisch 1982). We found only a very small correlation between reported age at menarche and age at the time of report. Whether this reflects a systematic telescoping effect, an actual decrease in age at menarche in the Dutch population or a combination of both, is uncertain.

Our results showed an estimate of 70% for the amount of total variance accounted for by additive genetic variation. Since in the analysis gene-environment interaction was ignored, its variance is attributed to non-shared variance. One may therefore conclude that between 70 and 71.5% of the total variance can be attributed to the effects of genotype (Falconer and Mackay 1996). This study showed high stability of menarche reports and relatively high heritability. This, coupled with results from other studies, suggests that recalled age at menarche is a fairly reliable trait that can be used in linkage and association studies.

Acknowledgments This study was funded by grants from the Netherlands Organisation for Scientific Research (NWO 575-25-006 and NWO 051-02-060, 668-772).

References

- Boomsma DI, Vink JM, Van Beijsterveldt TC et al (2002) Netherlands twin register: a focus on longitudinal research. *Twin Res* 5:401–406
- Ebling FJP (2005) The neuroendocrine timing of puberty. *Reproduction* 129:675–683
- Falconer DS, Mackay TFC (1996) Introduction to quantitative genetics, 4th edn. Longman, Harlow
- Greif EB, Ulman KJ (1982) The psychological impact of menarche on early adolescent females: a review of the literature. *Child Dev* 53:1413–1430
- Ibanez L, Ferrer A, Marcos MV, Hierro FR, de Zegher F (2000) Early puberty: rapid progression and reduced final height in girls with low birth weight. *Pediatrics* 106:e72
- Jinks JL, Fulker DW (1970) Comparison of the biometrical, MAVA, and classical approaches to the analysis of human behavior. *Psychol Bull* 73:311–349
- Kaprio J, Rimpela A, Winter T, Viken RJ, Rimpela M, Rose RJ (1995) Common genetic influences on BMI and age at menarche. *Hum Biol* 67:739–753
- Koopmans JR, Slutske WS, Heath A, Neale MC, Boomsma DI (1999) The genetics of smoking initiation and quantity smoked in Dutch adolescents and young adult twins. *Behav Genet* 29:383–393
- Loesch DZ, Hopper JL, Rogucka E, Huggins RM (1995) Timing and genetic rapport between growth in skeletal maturity and height around puberty: similarities and differences between girls and boys. *Am J Hum Genet* 56:753–759
- Meyer JM, Eaves LJ, Heath AC, Martin NG (1991) Estimating genetic influences on the age-at-menarche: a survival analysis approach. *Am J Med Genet* 39:148–154
- Must A, Phillips SM, Naumova EN, Blum M, Harris S, Dawson-Hughes B et al (2002) Recall of early menstrual history and

- menarcheal body size: after 30 years, how well do women remember? *Am J Epidemiol* 155:672–679
- Ojeda SR, Lomniczi A, Mastronardi C, Heger S, Roth C, Parent AS et al (2006) Minireview: the neuroendocrine regulation of puberty: is the time ripe for a systems biology approach? *Endocrinology* 147:1166–1174
- Palmert MR, Hirschhorn JN (2003) Genetic approaches to stature, pubertal timing, and other complex traits. *Mol Genet Metab* 80:1–10
- Pickles A, Pickering K, Simonoff E, Silberg J, Meyer J, Maes H (1998) Genetic "clocks" and "soft" events: a twin model for pubertal development and other recalled sequences of developmental milestones, transitions, or ages at onset. *Behav Genet* 28:243–253
- Pillemer DB, Koff E, Rhinehart ED, Rierdan J (1987) Flashbulb memories of menarche and adult menstrual distress. *J Adolesc* 10:187–199
- Schmidt FL, Hunter JE (1996) Measurement error in psychological research: lessons from 26 research scenarios. *Psychol Methods* 1:199–223
- Sisk CL, Foster DL (2004) The neural basis of puberty and adolescence. *Nat Neurosci* 7:1040–1047
- Stubbe JH, Posthuma D, Boomsma DI, de Geus EJC (2005) Heritability of life satisfaction in adults: a twin-family study. *Psychol Med* 35:1581–1588
- Towne B, Czerwinski S, Demerath EW, Blangero J, Roche AF, Siervogel RM (2005) Heritability of age at menarche in girls from the Fels longitudinal study. *Am J Phys Anthropol* 128:210–219
- Treloar SA, Martin NG (1990) Age at menarche as a fitness trait: nonadditive genetic variance detected in a large twin sample. *Am J Hum Genet* 47:137–148
- van den Berg SM, Setiawan A, Bartels M, Polderman TJC, Van der Vaart AW, Boomsma DI (2006) Individual differences in puberty onset in girls: Bayesian estimation of heritabilities and genetic correlations. *Behav Genet* 36:261–270
- Wink JM, Willemsen G, Stubbe JH, Middeldorp CM, Ligthart RSL, Baas K et al (2004) Estimating non-response bias in family studies: application to mental health and lifestyle. *Eur J Personal* 19:623–630
- Willemsen G, Posthuma D, Boomsma D (2005) Environmental factors determine where the Dutch live: results from the Netherlands Twin Register. *Twin Res Hum Genet* 8:312–317
- Wyshak G, Frisch RE (1982) Evidence for a secular trend in age of menarche. *N Engl J Med* 306:1033–1035
- Zacharias L, Wurtman RJ, Schatzoff M (1970) Sexual maturation in contemporary American girls. *Am J Obstet Gynecol* 108:833–846