

ORIGINAL ARTICLE

The *SNAP-25* gene is associated with cognitive ability: evidence from a family-based study in two independent Dutch cohorts

MF Gosso^{1,2,3}, EJC de Geus^{1,3}, MJ van Belzen^{2,3}, TJC Polderman^{1,4}, P Heutink^{1,2,3}, DI Boomsma^{1,2,3} and D Posthuma^{1,3}

¹Department of Biological Psychology, Vrije Universiteit, Amsterdam, The Netherlands; ²Section of Medical Genomics, Department of Clinical Genetics and Anthropogenetics, VU Medical Center, Amsterdam, The Netherlands; ³Center for Neurogenomics and Cognitive Research – CNCR, Vrije Universiteit, Amsterdam, The Netherlands and ⁴Department of Child and Adolescent Psychiatry, Erasmus University, Rotterdam, The Netherlands

The synaptosomal-associated protein of 25 kDa (*SNAP-25*) gene plays an integral role in synaptic transmission, and is differentially expressed in the mammalian brain in the neocortex, hippocampus, anterior thalamic nuclei, substantia nigra and cerebellar granular cells. Recent studies have suggested a possible involvement of *SNAP-25* in learning and memory, both of which are key components of human intelligence. In addition, the *SNAP-25* gene lies in a linkage area implicated previously in human intelligence. In two independent family-based Dutch samples of 391 (mean age 12.4 years) and 276 (mean age 37.3 years) subjects, respectively, we genotyped 12 single-nucleotide polymorphisms (SNPs) in the *SNAP-25* gene on 20p12–20p11.2. From all individuals, standardized intelligence measures were available. Using a family-based association test, a strong association was found between three SNPs in the *SNAP-25* gene and intelligence, two of which showed association in both independent samples. The strongest, replicated association was found between SNP rs363050 and performance IQ (PIQ), where the A allele was associated with an increase of 2.84 PIQ points ($P=0.0002$). Variance in this SNP accounts for 3.4 % of the phenotypic variance in PIQ.

Molecular Psychiatry (2006) 11, 878–886. doi:10.1038/sj.mp.4001868; published online 27 June 2006

Keywords: genetics; intelligence; synaptosomal; family-based; candidate gene approach; haplotype analysis

Introduction

Intelligence is one of the most heritable traits in humans, with heritability estimates ranging from 25 to 40% in early childhood¹ to 80% in adulthood.² Recently, the first genome-wide scan for intelligence was published, identifying two regions on chromosome 2q and 6p that showed significant linkage to intelligence, and several other regions showing suggestive linkage (4p, 7q, 20p, 21p).³ Other scans followed shortly, replicating the 6p region, and also pointing to other regions (e.g. 14q).^{4–7} An alternative approach to gene finding is to perform genetic association tests with candidate genes that are selected based on prior knowledge of biochemical functioning. We followed the latter approach and selected a putative candidate gene that was recently shown to be involved in learning and memory, which

are two major components of intelligence. Several studies have demonstrated that the hippocampus plays a central role in learning and memory.^{4–8} Damage to the hippocampus selectively impairs the ability to learn and remember.^{8–15} The synaptosomal-associated protein of 25 kDa (*SNAP-25*) gene lies in an area of previous suggestive linkage to intelligence (20p12–p11.2),³ and is highly expressed by neurons in the hippocampus.^{16–18} The *SNAP-25* gene product is a presynaptic plasma membrane protein that is an integral component of the vesicle docking and fusion machinery that regulates neurotransmitter release.^{17,19,20} It is also implicated in axonal growth and synaptic plasticity.²¹ Three lines of evidence suggest a major role of *SNAP-25* in learning and memory in humans. Firstly, selective inhibition of *SNAP-25* expression prevents axonal elongation and the transformation of growth cones to synaptic terminals,²¹ especially in hippocampal neurons.²² Such remodeling of nerve terminals in the adult brain may serve as a morphological substrate of learning and memory.^{21,23} Secondly, mRNA levels of *SNAP-25* are increased after the induction of long-term potentiation (LTP) in granule cells of the dentate gyrus.²⁴

Correspondence: Dr MF Gosso, Department of Biological Psychology, Vrije Universiteit, Van der Boecheorststraat 1, 1081 BT Amsterdam, The Netherlands.

E-mail: mf.gosso@vumc.nl

Received 3 March 2006; revised 31 May 2006; accepted 5 June 2006; published online 27 June 2006

Hippocampal LTP is thought to be a form of synaptic plasticity that underlies memory and learning.^{25–28} Thirdly, inhibition of hippocampal SNAP-25 leads to impaired long-term contextual fear memory, spatial memory, as well as decreased LTP.²³ The suggestive (according to the Lander and Kruglyak guidelines)²⁹ linkage finding of general intelligence (20p12–p11.2)³ to the area containing SNAP-25 renders this gene a putative candidate gene for human intelligence.

The present study aims to investigate whether SNAP-25 gene plays a role in human intelligence. To this end, a family-based association approach is used in two independent cohorts of children (mean age 12.4 years) and adults (mean age 37.3 years).

Materials and methods

Subjects

All twins and their siblings were part of two larger cognitive studies and were recruited from the Netherlands Twin Registry.³⁰ Informed consent was obtained from the participants (adult cohort) or from their parents if they were under 18 (young cohort). The current study was approved by the institutional review board of the VU University Medical Center. None of the individuals tested suffered from severe physical or mental handicaps, as assessed through standard questionnaire.

Young cohort

The young cohort consisted of 177 twin pairs born between 1990 and 1992, and 55 siblings.³¹ The twins were 12 (mean = 12.4, s.d. = 0.95) years of age and the siblings were between 8 and 15 years old at the time of testing. There were 41 monozygotic male twin pairs (MZM), 28 dizygotic male twin pairs (DZM), 56 monozygotic female twin pairs (MZF), 25 dizygotic female twin pairs (DZF), 27 dizygotic opposite-sex twin pairs (DOS), 28 male siblings and 27 female siblings. Participation in this study included a voluntary agreement to provide buccal swabs for DNA extraction.

Adult cohort

A total of 793 family members from 317 extended twin families participated in the adult cognition

study.² Participation in this study did not automatically include DNA collection; however, part of the sample (276 subjects) returned to the lab to provide blood for DNA extraction. Mean age was 37.3 years (s.d. = 12.50). There were 20 MZM, 11 DZM, one DZM triplet, 14 MZF, 22 DZF and 17 DOS, 23 female siblings and 23 male siblings, and 59 subjects from incomplete twin pairs (18 males, 41 females).

Cognitive testing

In the young cohort, cognitive ability was assessed with the Dutch adaptation of the Wechsler Intelligence Scale for Children-Revised,³² and consisted of four verbal subtests (similarities, vocabulary, arithmetic and digit span) and two performance subtests (block design and object assembly).

In the adult cohort, the Dutch adaptation of the Wechsler Adult Intelligence Scale III-Revised³³ assessed IQ and consisted of four verbal subtests (information, similarities, vocabulary and arithmetic) and four performance subtests (picture completion, block design, matrix reasoning and digit-symbol substitution). In both cohorts, verbal IQ (VIQ), performance IQ (PIQ) and full-scale IQ (FSIQ) were normally distributed. Correlations between FSIQ/VIQ, FSIQ/PIQ and PIQ/VIQ were 0.89, 0.81 and 0.45, respectively, in the young cohort, and 0.90, 0.84 and 0.55, respectively, in the adult cohort. Means and standard deviations of the full and genotyped cohorts are provided in Table 1.

DNA collection and genotyping

Buccal swabs were obtained from 391 children; blood was obtained from 276 adults. The DNA isolation from buccal swabs was performed using a chloroform/isopropanol extraction.³⁴ DNA was extracted from blood samples using the salting out protocol.³⁵

Zygosity was assessed using 11 polymorphic microsatellite markers (Het > 0.80). Tagging single-nucleotide polymorphisms (*tag*-SNPs) selection criteria were defined as SNPs with a minor allele frequency (MAF) above 0.10 and genotypic correlation (ρ) across the genotypes of maximal 0.85 as obtained from a randomly selected Caucasian sample (http://www.celeradiagnostics.com/cdx/applera_genomics). MAF had to be > 0.10 in order to avoid the rare heterozygous genotypes and SNPs with a ρ above

Table 1 Means and standard deviations of PIQ, VIQ and FSIQ in the young and adult cohorts

N	Young cohort		Adult cohort	
	Total sample 409	Genotyped 391	Total sample 793	Genotyped 276
Age (s.d.)	12.37 (0.95)	12.36 (0.90)	37.60 (13.00)	37.40 (12.42)
Mean PIQ (s.d.)	101.40 (12.85)	101.66 (12.96)	100.96 (12.50)	100.04 (12.40)
Mean VIQ (s.d.)	98.42 (19.04)	98.90 (19.02)	92.78 (13.83)	93.03 (14.36)
Mean FSIQ (s.d.)	99.81 (15.20)	100.21 (15.21)	95.74 (11.62)	95.59 (12.04)

Abbreviations: FSIQ, full-scale IQ; PIQ, performance IQ; VIQ, verbal IQ.




Figure 1 Location of *tag*-SNPs selected within the *SNAP-25* gene on chromosome 20 p12–p11.2.

0.85 with any of the other SNPs were not selected, to avoid redundancy. Twelve *tag*-SNPs in the *SNAP-25* gene were selected according to these criteria (<http://www.appliedbiosystems.com/support/software/snplex/>) using SNP Browser version 2.0.4 (NCBI build 34). Ranging from the 5' untranslated region (5' UTR) to 3'UTR region within the *SNAP-25* gene, the following SNPs were selected: rs883381, rs1889189, rs363039, rs363050, rs362569, rs6039806, rs362990, rs1051312, rs8636, rs362602, rs362552 and rs725919 (see Figure 1). Genotyping was performed blind to familial status and phenotypic data. Both MZ twins of a pair were included in genotyping, serving as additional controls.

The SNplex assay was conducted following the manufacturer's recommendations (Applied Biosystems, Foster city, CA, USA). All pre-PCR steps were performed on a cooled block. Reactions were carried out in Gene Amp 9700 Thermocycler (Applied Biosystems, Foster city, CA, USA). PCR products were analyzed with ABI3730 Sequencer (Applied Biosystems, Foster city, CA, USA). Data were analyzed using Genemapper v3.7 (Applied Biosystems, Foster city, CA, USA).

Statistical analyses

Allele frequencies of the 12 selected *tag*-SNPs were estimated in both young and adult cohorts using Pedstats (<http://www.sph.umich.edu/csg/abecasis/PedStats>) in which a Hardy–Weinberg test is implemented, based on an exact calculation of the probability of observing a certain number of heterozygotes conditional on the number of copies of the minor SNP allele. MZ twins were considered as one genotype, when estimating allele frequencies.

Linkage disequilibrium (LD) parameters (D' and r^2) were calculated from the haplotype frequency estimates using Haploview 3.2 (<http://www.broad.mit.edu/mpg/haploview>). $D' = 1$ if, and only if, two SNPs have not been separated by recombination (or recurrent mutation). This LD parameter is sensitive to sample size, especially when SNPs with rare allele frequencies are considered. The value of $r^2 = 1$ if, and only if, the SNPs have not been separated by recombination and have the same allele frequency. For quantifying and comparing LD in the context of mapping, r^2 is slightly preferred.³⁶ Values of r^2 ranged from 0.001 to 0.680 in our sample, conforming relatively low LD between the separate *tag*-SNPs (see Table 2).

Haplotypes were estimated using SNPs that showed a significant association with IQ in both samples, using the expectation-maximization (EM) algorithm to obtain the maximum likelihood estimates of haplotype frequencies in each sample,³⁷ as implemented in the Allegro software package.³⁸ The EM algorithm allows for missing data and can be applied when no parental genotypes are available.

Genetic association tests were conducted using the program QTDT, which implements the orthogonal association model proposed by Abecasis *et al.*³⁹ (see also Fulker *et al.*;⁴⁰ extended by Posthuma *et al.*⁴¹). This model allows the decomposition of the genotypic association effect into orthogonal between- (β_b) and within- (β_w) family components, can incorporate fixed effects of covariates and can also model the residual sib-correlation as a function of polygenic or environmental factors. MZ twins can be included and are modeled as such, by adding zygosity status to the datafile. They are not informative to the within-family association component (unless they are paired with non-twin siblings), but are informative for the between-family component. The between-family association component is sensitive to population admixture, whereas the within-family component is significant only in the presence of LD owing to close linkage. If population stratification acts to create a false association, the test for association using the within-family component is still valid, and provides a conservative test of association. Testing for the equality of the β_b and β_w effects serves as a test of population stratification. If this test is not significant, the between- and within-family effects are equal and total association test that uses the whole population at once can be applied. It should be noted, however, that given the relatively modest sample size, both the within-family test and the population stratification test are not as powerful as the 'total' association test. As we tested multiple SNPs, a significance level of 0.01 was kept.

Results

Single SNP analysis

In total, 391 subjects for the young cohort and 276 subjects for the adult cohort were available for SNP genotyping. Based on blind controls and MZ checks, no genotyping errors were found. Eight SNPs out of the 12 selected were in Hardy–Weinberg equilibrium (HWE) in both cohorts. SNPs not in HWE (rs362990,

Table 2 Estimates of LD parameters r^2 (lower) and D' (upper) for tag-SNPs within the SNAP-25 gene

	rs883381	rs1889189	rs363039	rs363050	rs362569	rs6039806	rs362990	rs1051312	rs8636	rs362602	rs362552	rs725919
rs883381	—											
rs1889189	0.223	0.885	0.495	0.491	0.012	0.014	0.144	0.154	0.021	0.428	0.149	0.133
rs363039	0.211	0.242	1.000	0.467	0.641	0.491	0.428	0.173	0.116	0.342	0.268	0.326
rs363050	0.192	0.078	—	0.828	0.072	0.071	0.195	0.184	0.066	0.316	0.177	0.234
rs362569	0.000	0.124	0.004	0.039	0.276	0.146	0.059	0.171	0.318	0.265	0.017	0.135
rs6039806	0.000	0.098	0.002	0.015	—	0.909	0.938	0.078	0.453	0.036	0.234	0.177
rs362990	0.012	0.031	0.027	0.001	0.630	—	0.971	0.099	0.394	0.037	0.248	0.165
rs1051312	0.009	0.019	0.012	0.014	0.486	0.397	—	0.226	0.536	0.122	0.230	0.187
rs8636	0.000	0.010	0.004	0.074	0.003	0.006	0.013	—	0.095	0.710	0.900	0.917
rs362602	0.070	0.085	0.033	0.034	0.078	0.079	0.062	0.002	—	0.098	0.811	0.888
rs362552	0.015	0.013	0.024	0.000	0.001	0.001	0.003	0.496	0.003	—	1.000	0.936
rs725919	0.008	0.014	0.030	0.0070	0.032	0.028	0.048	0.240	0.155	0.258	—	0.977
					0.013	0.009	0.027	0.176	0.134	0.158	0.682	—

Abbreviations: LD, linkage disequilibrium; SNAP-25, synaptosomal-associated protein of 25 kDa; tag-SNP, tagging single-nucleotide polymorphism.

rs6039806, rs362569 and rs1051312) were not included in further analyses. SNP rs883381 had a success rate of 80% in the young cohort; for all other SNPs in HWE, success rates were between 96.0 and 98.0% (see Table 3).

The models used in QTDT included effects of age and sex on the means and modeled additive allelic between- and within-family effects. Residual sib-correlations were modeled as a function of polygenic additive effects and non-shared environmental effects. Tests for the presence of population stratification were all nonsignificant, indicating that genotypic effects within families were not significantly different from those observed between families, suggesting that the more powerful total association test can be interpreted. Three SNPs (rs363039, rs363050, rs362602) showed significant associations with IQ. Two of these SNPs (rs363039, rs363050) were associated with IQ in both the young cohort and the independent adult cohort, showing association in the same direction and the same order of magnitude. The third SNP (rs362602) was seen as a trend to significant association only in the adult cohort. When we combined the two cohorts, the strongest association was seen between PIQ and rs363050, which is located on the 5'UTR of the SNAP-25 gene ($\chi^2 = 13.56$, $P = 0.0002$). The increaser allele of this SNP was associated with an increase of 2.84 IQ points (see Tables 4 and 5 and Figure 2).

Within-family association tests are based on all siblings that are part of pairs with contrasting genotypes within a family and are thus less powerful than total association tests. The latter is preferred if there is no evidence of population stratification. It is, however, interesting to check whether the significant associations observed in the total association test are also present when looking only at the within-family association. In the within-association test, for SNP rs363039, a trend was seen in both cohorts separately, whereas the G allele was suggestive of association ($P < 0.05$) in the combined cohort. For SNP rs363050, the within-family association with the A allele was suggestive ($P = 0.06$) in the combined cohort. SNPs rs8636 and rs362602 were significant in the adult cohort ($P < 0.01$) when only considering the within-family test. These results support the results as found using the more powerful total association test.

Haplotype analysis

The two SNPs that showed a significant association with IQ in both cohorts were 13 kb apart. Because these SNPs are in LD with each other ($r^2 = 0.46$), these SNPs were used to estimate haplotypes within each sample. Haplotype analysis of SNPs that are in LD with each other is more powerful than single SNP analysis because the combination of SNPs into a haplotype can be considered as a multiallelic marker that is more informative than a biallelic marker. Nonsignificant SNPs were not used for further haplotype analysis, as all SNPs were selected on the basis of being tagging SNPs. From Table 2, it can be

Table 3 List of selected *tag*-SNP within the *SNAP*-25 gene with their estimated heterozygosity rates for the young/adult cohort

Name	Position	Gene location	Obs HET	Pred HET	Success rate	MAF	HWE
rs883381	10160727	5'UTR	0.41/0.54	0.47/0.46	80.10/94.80	0.38/0.37	OK/OK
rs1889189	10192086	5'UTR	0.40/0.37	0.43/0.44	100.00/99.30	0.32/0.33	OK/OK
rs363039	10215496	Intron 1	0.42/0.50	0.42/0.47	99.00/100.00	0.30/0.37	OK/OK
rs363050	10229257	Intron 1	0.52/0.47	0.49/0.50	100.00/100.00	0.42/0.47	OK/OK
<i>rs362569</i>	<i>10241733</i>	<i>Intron 1</i>	<i>0.37/0.50</i>	<i>0.46/0.48</i>	<i>90.70/91.00</i>	<i>0.36/0.41</i>	<i>Not in HWE/OK</i>
<i>rs6039806</i>	<i>10253654</i>	<i>Intron 3</i>	<i>0.38/0.49</i>	<i>0.50/0.50</i>	<i>92.80/88.00</i>	<i>0.46/0.47</i>	<i>Not in HWE/OK</i>
<i>rs362990</i>	<i>10271221</i>	<i>Intron 5</i>	<i>0.30/0.41</i>	<i>0.39/0.40</i>	<i>100.00/96.30</i>	<i>0.27/0.27</i>	<i>Not in HWE/OK</i>
<i>rs1051312</i>	<i>10282088</i>	<i>Exon 8</i>	<i>0.22/0.21</i>	<i>0.46/0.47</i>	<i>77.80/71.50</i>	<i>0.37/0.38</i>	<i>Not in HWE/Not in HWE</i>
rs8636	10282742	Exon 8	0.44/0.51	0.45/0.49	98.20/98.90	0.34/0.42	OK/OK
rs362602	10288528	3'UTR	0.42/0.47	0.47/0.47	100.00/98.50	0.39/0.38	OK/OK
rs362552	10291217	3'UTR	0.42/0.39	0.43/0.39	99.20/97.40	0.29/0.26	OK/OK
rs725919	10298094	3'UTR	0.34/0.34	0.36/0.33	99.00/98.90	0.23/0.21	OK/OK

Abbreviations: HWE, Hardy–Weinberg equilibrium; MAF, minor allele frequency; Obs HET, observed heterozygosity; Pred HET, predicted heterozygosity; *SNAP*-25, synaptosomal-associated protein of 25 kDa; *tag*-SNP, tagging single-nucleotide polymorphism; UTR, untranslated region.

Note: *Tag*-SNPs were selected if allele frequency was >10% (18.0% heterozygosity) and a genotypic correlation (ρ) <0.85. SNPs not in HWE are in italics.

Table 4 Family-based association analysis for *SNAP*-25 *tag*-SNPs for young, adult and combined cohort

<i>tag</i> -SNP position (bp)	Phenotype	Young cohort				Adult cohort				Combined			
		N	χ^2	P	Genotypic effect	N	χ^2	P	Genotypic effect	N	χ^2	P	Genotypic effect
rs883381 (10160727)	PIQ	308	2.05	0.152	1.59 (G)	253	0.95	0.329	1.24 (G)	561	3.07	0.080	1.46 (G)
	VIQ	306	0.00	0.981	0.04 (T)	254	2.24	0.134	2.00 (G)	560	0.55	0.457	0.80 (G)
	FSIQ	308	0.30	0.584	0.70 (G)	252	1.88	0.170	1.62 (G)	560	1.50	0.221	1.09 (G)
rs1889189 (10192086)	PIQ	385	0.07	0.788	0.30 (T)	268	1.13	0.287	1.22 (T)	653	0.63	0.427	0.63 (T)
	VIQ	383	2.54	0.111	2.56 (C)	269	1.88	0.170	1.68 (T)	652	0.59	0.444	0.81 (C)
	FSIQ	385	0.94	0.332	1.25 (T)	267	1.91	0.166	1.49 (T)	652	0.03	0.861	0.15 (C)
rs363039 (10215496)	PIQ	381	6.52	0.010	2.99 (G)	271	3.28	0.070	2.08 (G)	652	9.21	0.002	2.51 (G)
	VIQ	379	3.21	0.073	3.09 (G)	272	6.37	0.012	3.14 (G)	651	7.88	0.005	3.12 (G)
	FSIQ	381	5.83	0.016	3.34 (G)	270	6.18	0.013	2.68 (G)	651	10.88	0.001	2.98 (G)
rs363050 (10229257)	PIQ	385	7.88	0.005	3.01 (A)	267	5.44	0.020	2.55 (A)	652	13.56	0.0002	2.84 (A)
	VIQ	383	1.94	0.164	2.21 (A)	268	6.03	0.014	2.89 (A)	651	5.91	0.015	2.52 (A)
	FSIQ	385	5.47	0.019	2.96 (A)	266	6.90	0.009	2.69(A)	651	11.48	0.0007	2.86 (A)
rs8636 (10282742)	PIQ	378	0.21	0.643	0.50 (T)	264	2.95	0.086	1.99 (C)	642	2.22	0.137	1.18 (T)
	VIQ	376	0.30	0.587	0.86 (T)	265	1.92	0.166	1.71 (C)	641	0.07	0.785	0.29 (C)
	FSIQ	378	0.00	0.978	0.03 (T)	261	3.39	0.065	1.99 (C)	641	1.08	0.299	0.90 (C)
rs362602 (10288528)	PIQ	385	0.33	0.567	0.59 (A)	265	3.01	0.082	1.97(G)	650	0.30	0.583	0.42 (G)
	VIQ	383	1.40	0.236	1.80 (A)	266	7.91	0.005	3.49 (G)	649	0.06	0.811	0.24 (G)
	FSIQ	385	1.16	0.281	1.31 (A)	264	7.57	0.006	2.96 (G)	649	0.22	0.639	0.40 (G)
rs362552 (10291217)	PIQ	382	1.01	0.314	1.09 (G)	260	0.30	0.587	0.73 (A)	642	0.32	0.572	0.47 (G)
	VIQ	380	0.31	0.575	0.90 (G)	261	1.20	0.273	1.59 (A)	641	0.01	0.924	0.10 (G)
	FSIQ	382	0.67	0.415	1.04 (G)	259	1.01	0.315	1.27 (A)	641	0.04	0.832	0.19 (G)
rs725919 (10298094)	PIQ	381	1.87	0.172	1.63 (A)	267	0.45	0.505	0.94 (G)	648	0.60	0.439	0.70 (A)
	VIQ	379	0.14	0.712	0.65 (A)	268	1.21	0.271	1.70 (G)	647	0.55	0.457	0.26 (A)
	FSIQ	381	0.65	0.420	1.13 (A)	266	1.11	0.292	1.40 (G)	647	0.05	0.831	0.21 (A)

Abbreviations: FSIQ, full-scale IQ; PIQ, performance IQ; *SNAP*-25, synaptosomal-associated protein of 25 kDa; *tag*-SNP, tagging single-nucleotide polymorphism; VIQ, verbal IQ.

Note: The genotypic effect is the increase in IQ points associated with the increaser allele. *P* values below <0.01 are in bold. Sex and age were included as covariates; residual variance was modeled as a function of polygenic effects and non-shared environmental effects.

Table 5 Means (s.d.) per genotype for PIQ, VIQ and FSIQ for young and adult cohorts in the four *tag*-SNPs within the *SNAP-25* gene that show association with a significant association

tag-SNP position (bp)	Young cohort					Adult cohort				
	Phenotype	Genotype			Total N	Genotype frequency			Total N	
rs363039 (10168496)	<i>Frequency</i>	GG 0.49	AG 0.42	AA 0.09		GG 0.38	AG 0.50	AA 0.12		
	Mean PIQ (s.d.)	103.68 (13.72)	100.58 (12.30)	97.06 (10.79)	381	101.57 (12.46)	96.89 (11.69)	98.96 (13.71)	271	
	Mean VIQ (s.d.)	101.13 (19.28)	97.22 (19.14)	97.79 (16.19)	379	95.53 (14.86)	90.62 (13.66)	89.71 (14.15)	272	
	Mean FSIQ (s.d.)	102.45 (16.09)	98.47 (14.63)	96.94 (13.54)	381	97.73 (12.66)	92.98 (11.28)	93.21 (13.27)	270	
	<i>Frequency</i>	AA 0.32	AG 0.52	GG 0.16		AA 0.29	AG 0.47	GG 0.24		
	Mean PIQ (s.d.)	104.26 (13.26)	101.11 (13.08)	98.24 (11.29)	385	102.50 (13.58)	97.60 (11.95)	97.34 (11.01)	267	
rs363050 (10182257)	Mean VIQ (s.d.)	103.12 (18.82)	96.52 (19.08)	98.68 (17.92)	383	95.26 (15.65)	92.26 (13.05)	89.20 (15.12)	268	
	Mean FSIQ (s.d.)	103.98 (15.33)	98.36 (15.44)	98.03 (13.92)	385	98.08 (13.72)	93.98 (11.42)	92.71 (11.45)	266	
	<i>Frequency</i>	AA 0.40	AG 0.42	GG 0.18		AA 0.39	AG 0.47	GG 0.14		
rs362602 (10241528)	Mean PIQ (s.d.)	102.45 (11.74)	101.45 (13.11)	100.44 (15.34)	385	100.23 (12.45)	99.54 (12.82)	100.11 (11.06)	265	
	Mean VIQ (s.d.)	99.97 (16.86)	99.50 (19.62)	95.82 (21.78)	383	92.45 (14.98)	91.11 (14.40)	97.63 (12.40)	266	
	Mean FSIQ (s.d.)	101.28 (12.81)	100.37 (15.77)	97.10 (19.01)	385	94.91 (12.59)	94.08 (12.43)	98.16 (10.99)	264	

Abbreviations: FSIQ, full-scale IQ; MZ, monozygotic; PIQ, performance IQ; QTDT, quantitative transmission disequilibrium test; *SNAP-25*, synaptosomal-associated protein of 25 kDa; *tag*-SNP, tagging single-nucleotide polymorphism; VIQ, verbal IQ.

Note: N denotes the number of individuals.




Figure 2 IQ means and standard error for the means for FSIQ, VIQ and PIQ for the combined cohort are plotted against the two most significant replicated tag-SNPs rs363050 and rs363039 genotypes.

seen that indeed LD among the tag-SNPs flanking the two most significant SNPs is very low, which would also be expected given the lack of association with these flanking SNPs and IQ.

Haplotypes were estimated using the two SNPs rs363039 and rs363050 that were associated with psychometric IQ scores. Four possible haplotypes were G-A, A-G, G-G and A-A, with haplotype frequencies 0.55, 0.29, 0.13 and 0.03, respectively, in the young cohort and 0.54, 0.31, 0.13 and 0.03, respectively, in the adult sample. Significant associations were found in both samples. When the data were combined, highly significant associations were observed with the G-A haplotype with FSIQ ($\chi^2(1) = 11.14, P = 0.0008$), VIQ ($\chi^2(1) = 7.15, P = 0.0074$) and PIQ ($\chi^2(1) = 10.61, P = 0.0011$) (see Table 6). These results confirm the single SNP association results.

Discussion

To investigate the possible role of the SNAP-25 gene in intelligence, we employed a family-based genetic association test in two independent cohorts of 391 children (mean age 12.4 years) and 276 adults (mean age 37.3 years). Replicated association was found in the two cohorts for two SNPs in the SNAP-25 gene. Strongest evidence was found for SNP rs363050 in intron 1 at the 5'UTR, showing an effect size of 2.84 IQ points ($P = 0.0002$) for the increaser allele. Haplotype analyses confirmed the region containing these two SNPs to be strongly associated with IQ.

Table 6 Family-based association analysis for SNAP-25 tagging haplotype for young, adult and combined cohort

Haplotype	Haplotype frequency		Phenotype			Young			Adult			Combined			
	Young	Adult	PIQ	VIQ	FSIQ	N	χ^2	P	N	χ^2	P	N	χ^2	P	Genotypic effect
G-A	0.55	0.54	PIQ			380	6.53	0.011	263	3.82	0.051	643	10.61	0.001	2.18
			VIQ	2.33		378	2.98	0.084	264	5.47	0.019	642	7.15	0.007	2.41
			FSIQ	2.64		380	5.91	0.015	262	5.75	0.016	642	11.14	0.001	2.44
A-G	0.29	0.31	PIQ	-1.98		380	2.54	0.111	263	4.19	0.041	643	6.45	0.011	-2.24
			VIQ	-1.65		378	0.82	0.365	264	5.63	0.018	642	4.24	0.040	-2.44
			FSIQ	-2.07		380	1.98	0.159	262	6.56	0.010	642	6.81	0.009	-2.53
G-G	0.13	0.13	PIQ	-1.48		380	1.23	0.267	263	0.17	0.680	643	1.49	0.222	-1.24
			VIQ	-0.79		378	0.16	0.689	264	0.09	0.764	642	0.23	0.632	-0.66
			FSIQ	-1.32		380	0.78	0.377	262	0.06	0.806	642	0.78	0.377	-0.98
A-A	0.03	0.03	PIQ	-1.97		380	1.27	0.260	263	0.07	0.791	643	1.14	0.286	-1.50
			VIQ	-3.74		378	2.13	0.144	264	0.58	0.446	642	2.65	0.104	-3.07
			FSIQ	-3.17		380	2.31	0.129	262	0.42	0.517	642	2.58	0.108	-2.48

Abbreviations: FSIQ, full-scale IQ; PIQ, performance IQ; SNAP-25, synaptosomal-associated protein of 25 kDa; VIQ, verbal IQ.

The *SNAP-25* gene, located on chromosome 20 p12–12p11.2, encodes a presynaptic terminal protein. In the mature brain, the *SNAP-25* gene product forms a complex with syntaxin and the synaptic vesicle proteins (synaptobrevin and synaptotagmin) that mediates exocytosis of neurotransmitter from the synaptic vesicle into the synaptic cleft (see Horikawa *et al.*,¹⁹ Seagar *et al.*,²⁰ Bark *et al.*,⁴² Low *et al.*⁴³). During development, *SNAP-25* is also involved in synaptogenesis, forming presynaptic sites and neuritic outgrowth.^{17,21} *SNAP-25* is thought to be differentially expressed in the brain, and is primarily present in the neocortex, hippocampus, anterior thalamic nuclei, substantia nigra and cerebellar granular cells. In the mature brain, expression is mainly seen at presynaptic terminals.¹⁷

SNAP-25 exists in two splicing variants in relation to exon 5, *SNAP-25a* and *SNAP-25b*. Both isoforms differ in only nine out of 39 amino acids encoded by the alternative spliced exons,⁴⁴ resulting in a differentiated membrane anchoring relative to cysteine residues involved in post-transcriptional fatty acylation.⁴⁵ Both isoforms are thought to be equally important but at different time points for both neuronal maturation and neurotransmitter release.^{21,22,42,46} Roberts *et al.*²⁴ demonstrated that mRNA levels of both isoforms are elevated after induction of LTP, suggesting a role of *SNAP-25* in synaptic plasticity. A recent study involving antisense oligonucleotides against *SNAP-25* at the hippocampal CA1 region reported the possible involvement of *SNAP-25* in learning and memory, particularly memory consolidation.²³ Steffensen *et al.*⁴⁷ found that hippocampal LTP is attenuated in hemizygous mice from the *Coloboma* mice strain. The *Coloboma* mice strain is characterized by a 2-cM deletion on the mouse homolog of chromosome 2, in a region containing the mouse *SNAP-25* gene. Mice hemizygous for this deletion exhibit a wide spectrum of phenotypic and neurological abnormalities such as ophthalmic deformation, head bobbing, circling, hyperactivity and small body size.^{45,48,49} Because of the observed increased hyperactivity of hemizygous *Coloboma* mice, the role of *SNAP-25* in attention deficit hyperactivity disorder (ADHD) has been tested in several studies.^{50–54} All, except one (Xu *et al.*⁵⁴), report a significant association of *SNAP-25* with ADHD in humans. The exact role of *SNAP-25* in ADHD, however, remains unknown. ADHD is a neuropsychiatric condition characterized by hyperactive behavior and impaired attentive ability, resulting in both social and academic dysfunction. The present study suggests that involvement of *SNAP-25* may not be specific to the hyperactivity component of ADHD, but plays a more general role in learning and memory, through its effect on LTP and synaptic plasticity.

Both individual and haplotype analyses were conducted with two SNPs that showed significant association with intelligence in our study, tagging the 5'UTR region of *SNAP-25* gene. Genetic (non)coding

variants lying within this non-coding region might be regulating this protein expression. These variants may influence regulatory binding sites, which in turn may modify gene expression and consequently neurotransmitter release regulation. Subtle changes in the fine-tuning at the neurotransmitter release machinery level, as well as in the interaction between neurotransmitter receptor subtypes, might be manifest in substantial differences when LTP is being achieved. This complex fine-tuning may be reflected as individual differences in memory and learning, two fundamental aspects of human intelligence. Future functional studies will provide the insight needed in order to disentangle the complex interplay among *SNAP-25* gene (non)coding variants and cognitive ability.

Acknowledgments

This study was supported by the Universitair Stimulerings Fonds (Grant no. 96/22), the Human Frontiers of Science Program (Grant no. rg0154/1998-B) and the Netherlands Organization for Scientific Research (NWO) Grants 904-57-94 and NWO/SPI 56-464-14192. DP was supported by GenomEUtwin Grant EU/QLRT-2001-01254 and by NWO/MaGW Vernieuwingsimpuls 016-065-318. This study was supported by the Centre for Medical Systems Biology (CMSB), a centre of excellence approved by the Netherlands Genomics Initiative/Netherlands Organization for Scientific Research (NWO). We thank Saskia van Mil and David Sondervan from the Medical Genomics Laboratory for technical support, and Zoltan Bochdanovits for valuable comments. We also like to thank the families from the Netherland Twin Registry (NTR) who participated in this study.

References

- 1 Bartels M, Rietveld M, Van Baal GCM, Boomsma DI. Genetic and environmental influences on the development of intelligence. *Behav Genet* 2002; **32**: 237–249.
- 2 McGue M, Bouchard Jr TJ, Iacono WG, Lykken DT. Behavioral genetics of cognitive ability: a life-span perspective. In: Plomin R, McClearn GE (eds). *Nature, Nurture, and Psychology*. American Psychological Association: Washington, DC, 1993, pp 59–76.
- 3 Posthuma D, Luciano M, Geus EJ, Wright MJ, Slagboom PE, Montgomery GW *et al*. A genomewide scan for intelligence identifies quantitative trait loci on 2q and 6p. *Am J Hum Genet* 2005; **77**: 318–326.
- 4 Buyske S, Bates M, Gharani N, Matise T, Tischfield J, Manowitz P. Cognitive traits link to human chromosomal regions. *Behav Genet* 2006; **36**: 65–76.
- 5 Dick D, Aliev F, Bierut L, Goate A, Rice J, Hinrichs A *et al*. Linkage analyses of IQ in the Collaborative Study on the Genetics of Alcoholism (COGA) sample. *Behav Genet* 2006; **36**: 77–86.
- 6 Luciano M, Wright MJ, Duffy DL, Wainwright MA, Zhu G, Evans DM *et al*. Genome-wide scan of IQ finds significant linkage to a quantitative trait locus on 2q. *Behavior Genetics* 2006. *Behav Genet* 2006; **36**: 45–55.
- 7 Wainwright M, Wright M, Luciano M, Montgomery G, Geffen G, Martin N. A linkage study of academic skills defined by the Queensland core skills test. *Behav Genet* 2006; **36**: 56–64.
- 8 Squire LR, Kandel ER (eds). *Memory: From Mind to Molecules*. Scientific American Library: New York, 1999.

- 9 Kim JJ, Rison RA, Fanselow MS. Effects of amygdala, hippocampus, and periaqueductal gray lesions on short- and long-term contextual fear. *Behav Neurosci* 1993; **107**: 1093–1098.
- 10 Logue SF, Paylor R, Wehner JM. Hippocampal lesions cause learning deficits in inbred mice in the Morris water maze and conditioned-fear task. *Behav Neurosci* 1997; **111**: 104–113.
- 11 Morris RG, Garrud P, Rawlins JN, O'Keefe J. Place navigation impaired in rats with hippocampal lesions. *Nature* 1982; **297**: 681–683.
- 12 Phillips RG, LeDoux JE. Lesions of the fornix but not the entorhinal or perirhinal cortex interfere with contextual fear conditioning. *J Neurosci* 1995; **15**: 5308–5315.
- 13 Sandin J, Ogren SO, Terenius L. Nociceptin/orphanin FQ modulates spatial learning via ORL-1 receptors in the dorsal hippocampus of the rat. *Brain Res* 2004; **997**: 222.
- 14 Selden NR, Everitt BJ, Jarrard LE, Robbins TW. Complementary roles for the amygdala and hippocampus in aversive conditioning to explicit and contextual cues. *Neuroscience* 1991; **42**: 335–350.
- 15 Sutherland RJ, Kolb B, Whishaw IQ. Spatial mapping: definitive disruption by hippocampal or medial frontal cortical damage in the rat. *Neurosci Lett* 1982; **31**: 271–276.
- 16 Geddes JW, Hess EJ, Hart RA, Kesslak JP, Cotman CW, Wilson MC. Lesions of hippocampal circuitry define synaptosomal-associated protein-25 (SNAP-25) as a novel presynaptic marker. *Neuroscience* 1990; **38**: 515–525.
- 17 Oyler GA, Higgins GA, Hart RA, Battenberg E, Billingsley M, Bloom FE *et al*. The identification of a novel synaptosomal-associated protein, SNAP-25, differentially expressed by neuronal subpopulations. *J Cell Biol* 1989; **109**: 3039–3052.
- 18 Frassoni C, Inverardi F, Coco S, Ortino B, Grumelli C, Pozzi D *et al*. Analysis of SNAP-25 immunoreactivity in hippocampal inhibitory neurons during development in culture and *in situ*. *Neuroscience* 2005; **131**: 813.
- 19 Horikawa HP, Saisu H, Ishizuka T, Sekine Y, Tsugita A, Odani S *et al*. A complex of rab3A, SNAP-25, VAMP/synaptobrevin-2 and syntaxins in brain presynaptic terminals. *FEBS Lett* 1993; **330**: 236–240.
- 20 Seagar M, Takahashi M. Interactions between presynaptic calcium channels and proteins implicated in synaptic vesicle trafficking and exocytosis. *J Bioenerg Biomembr* 1998; **30**: 347–356.
- 21 Osen-Sand A, Catsicas M, Staple JK, Jones KA, Ayala G, Knowles J *et al*. Inhibition of axonal growth by SNAP-25 antisense oligonucleotides *in vitro* and *in vivo*. *Nature* 1993; **364**: 445–448.
- 22 Grosse G, Grosse J, Tapp R, Kuchinke J, Gorsleben M, Fetter I *et al*. SNAP-25 requirement for dendritic growth of hippocampal neurons. *J Neurosci Res* 1999; **56**: 539–546.
- 23 Hou Q, Gao X, Zhang X, Kong L, Wang X, Bian W *et al*. SNAP-25 in hippocampal CA1 region is involved in memory consolidation. *Eur J Neurosci* 2004; **20**: 1593–1603.
- 24 Roberts LA, Morris BJ, O'Shaughnessy CT. Involvement of two isoforms of SNAP-25 in the expression of long-term potentiation in the rat hippocampus. *NeuroReport* 1998; **9**: 33–36.
- 25 Bliss TV, Collingridge GL. A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 1993; **361**: 31–39.
- 26 Martin SJ, Grimwood PD, Morris RG. Synaptic plasticity and memory: an evaluation of the hypothesis. *Annu Rev Neurosci* 2000; **23**: 649–711.
- 27 Morris RG. Synaptic plasticity and learning: selective impairment of learning rats and blockade of long-term potentiation *in vivo* by the N-methyl-D-aspartate receptor antagonist AP5. *J Neurosci* 1989; **9**: 3040–3057.
- 28 Morris RG, Anderson E, Lynch GS, Baudry M. Selective impairment of learning and blockade of long-term potentiation by an N-methyl-D-aspartate receptor antagonist, AP5. *Nature* 1986; **319**: 774–776.
- 29 Lander E, Kruglyak L. Genetic dissection of complex traits: guideline for interpreting and reporting linkage results. *Nat Genet* 1995; **11**: 241–247.
- 30 Boomsma DI. Twin registers in Europe: an overview. *Twin Res* 1998; **1**: 34–51.
- 31 Polderman TJC, Stins JF, Posthuma D, Gosso MF, Verhulst FC, Boomsma DI. The phenotypic and genotypic relation between working memory speed and capacity. Available online, 3 May 2006. doi:10.1016/J.intell.2006.03.010.
- 32 Wechsler D (ed). *Wechsler Intelligence Scale for Children-Revised (WISC-R)*. Swets & Zeitlinger: Lisse, 1986.
- 33 Wechsler D (ed). *WAIS-III Wechsler Adult Intelligence Scale*. Psychological Corporation: San Antonio, TX, 1997.
- 34 Meulenbelt I, Droog S, Trommelen GJ, Boomsma DI, Slagboom PE. High-yield noninvasive human genomic DNA isolation method for genetic studies in geographically dispersed families and populations. *Am J Hum Genet* 1995; **57**: 1252–1254.
- 35 Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988; **16**: 1215.
- 36 Ardlie KG, Kruglyak L, Seielstad M. Patterns of linkage disequilibrium in the human genome. *Nat Rev Genet* 2002; **3**: 299–309.
- 37 Excoffier L, Slatkin M. Maximum-likelihood estimation of molecular haplotype frequencies in a diploid population. *Mol Biol Evol* 1995; **12**: 921–927.
- 38 Gudbjartsson DF, Thorvaldsson T, Kong A, Gunnarsson G, Ingolfsdottir A. Allegro version 2. *Nat Genet* 2005; **37**: 1015–1016.
- 39 Abecasis GR, Cardon LR, Cookson WO. A general test of association for quantitative traits in nuclear families. *Am J Hum Genet* 2000; **66**: 279–292.
- 40 Fulker DW, Cherny SS, Sham PC, Hewitt JK. Combined linkage and association sib-pair analysis for quantitative traits. *Am J Hum Genet* 1999; **64**: 259–267.
- 41 Posthuma D, de Geus EJ, Boomsma DI, Neale MC. Combined linkage and association tests in mx. *Behav Genet* 2004; **34**: 179–196.
- 42 Bark IC, Wilson MC. Human cDNA clones encoding two different isoforms of the nerve terminal protein SNAP-25. *Gene* 1994; **139**: 291–292.
- 43 Low P, Norlin T, Risinger C, Larhammar D, Pieribone VA, Shupliakov O *et al*. Inhibition of neurotransmitter release in the lamprey reticulospinal synapse by antibody-mediated disruption of SNAP-25 function. *Eur J Cell Biol* 1999; **78**: 787–793.
- 44 Bark C, Wilson MC. Alternative splicing generates a variant SNAP-25 protein during development. *Soc Neurosci Abstr* 1991; **17**: 531.
- 45 Hess DT, Slater TM, Wilson MC, Skene JH. The 25 kDa synaptosomal-associated protein SNAP-25 is the major methionine-rich polypeptide in rapid axonal transport and a major substrate for palmitoylation in adult CNS. *J Neurosci* 1992; **12**: 4634–4641.
- 46 Bark IC, Hahn KM, Ryabinin AE, Wilson MC. Differential expression of SNAP-25 protein isoforms during divergent vesicle fusion events of neural development. *Proc Natl Acad Sci USA* 1995; **92**: 1510–1514.
- 47 Steffensen SC, Wilson MC, Henriksen SJ. Coloboma contiguous gene deletion encompassing Snap alters hippocampal plasticity. *Synapse* 1996; **22**: 281–289.
- 48 Heyser CJ, Wilson MC, Gold LH. Coloboma hyperactive mutant exhibits delayed neurobehavioral developmental milestones. *Brain Res Dev Brain Res* 1995; **89**: 264–269.
- 49 Snell GD, Bunker HP. Coloboma linkage. *Mouse News Lett* 1967; **37**: 34.
- 50 Brophy K, Hawi Z, Kirley A, Fitzgerald M, Gill M. Synaptosomal-associated protein 25 (SNAP-25) and attention deficit hyperactivity disorder (ADHD): evidence of linkage and association in the Irish population. *Mol Psychiatry* 2002; **7**: 913–917.
- 51 Feng Y, Crosbie J, Wigg K, Pathare T, Ickowicz A, Schachar R *et al*. The SNAP25 gene as a susceptibility gene contributing to attention-deficit hyperactivity disorder. *Mol Psychiatry* 2005; **10**: 998–1005.
- 52 Kustanovich V, Merriman B, McGough J, McCracken JT, Smalley SL, Nelson SF. Biased paternal transmission of SNAP-25 risk alleles in attention-deficit hyperactivity disorder. *Mol Psychiatry* 2003; **8**: 309–315.
- 53 Mill J, Curran S, Kent L, Gould A, Hockett L, Richards S *et al*. Association study of a SNAP-25 microsatellite and attention deficit hyperactivity disorder. *Am J Med Genet* 2002; **114**: 269–271.
- 54 Xu X, Knight J, Brookes K, Mill J, Sham P, Craig I *et al*. DNA pooling analysis of 21 norepinephrine transporter gene SNPs with attention deficit hyperactivity disorder: no evidence for association. *Am J Med Genet B Neuropsychiatr Genet* 2005; **134**: 115–118.